

POLARIZATION PHENOMENA IN FRAGMENTATION OF DEUTERONS TO PIONS AND NON-NUCLEON DEGREES OF FREEDOM IN THE DEUTERON

A.Yu. Illarionov[†], A.G. Litvinenko and G.I. Lykasov

Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia

[†] *E-mail: Alexei.Illarionov@jinr.ru*

Abstract

The fragmentation of deuterons into pions emitted forward in the kinematic region forbidden for free nucleon-nucleon collisions is analyzed. It is shown that the inclusion of the non-nucleon degrees of freedom in a deuteron allows to describe the experimental data about inclusive pion spectrum rather satisfactory and improves the description of data concerning the deuteron analyzing power T_{20} . The experimental data show the positive sign and very small values, less than 0.2, of T_{20} what can't be reproduced by the calculations ignoring these degrees of freedom.

Let us analyze the inclusive process $Dp \rightarrow \pi X$ for the polarized deuteron and pions emitted to forward at initial energies of order few GeV. If the deuteron is tensor analyzed and has p_D^{ZZ} component, then the inclusive spectrum of this reaction can be written in the form:

$$\rho_{pD}^{\pi}(p_D^{ZZ}) = \rho_{pD}^{\pi} [1 + A_{ZZ} \cdot p_D^{ZZ}] , \quad (1)$$

where $\rho_{pD}^{\pi} \equiv \varepsilon_{\pi} \cdot d\sigma_{pD}^{\pi}/d^3p_{\pi}$ is the inclusive spectrum for the case of unpolarized deuterons and $A_{ZZ} = \sqrt{2}T_{20}$ is the tensor analyzing power. They can be written in a fully covariant manner within the Bethe-Salpeter formalism [1]:

$$\rho_{pD}^{\pi} = \frac{1}{(2\pi)^3} \int \frac{\sqrt{\lambda(p, n)}}{\sqrt{\lambda(p, D)}} [\rho_{pN}^{\pi} \cdot \Phi^{(u)}(|\mathbf{q}|)] \frac{d^3q}{E_{\mathbf{q}}} ; \quad (2)$$

$$\rho_{pD}^{\pi} \cdot A_{ZZ} = -\frac{1}{(2\pi)^3} \int \frac{\sqrt{\lambda(p, n)}}{\sqrt{\lambda(p, D)}} [\rho_{pN}^{\pi} \cdot \Phi^{(t)}(|\mathbf{q}|)] \left(\frac{3 \cos^2 \theta_{\mathbf{q}} - 1}{2} \right) \frac{d^3q}{E_{\mathbf{q}}} \quad (3)$$

where $\lambda(p_1, p_2) \equiv (p_1 p_2)^2 - m_1^2 m_2^2 = \lambda(s_{12}, m_1^2, m_2^2)/4$ is the flux factor; p, n are the four-momenta of the proton-target and intra-deuteron nucleon, respectively; ρ_{pN}^{π} is the relativistic invariant inclusive spectrum of pions produced by interacting the intra-deuteron nucleon with the proton-target. The functions $\Phi^{(u)}(|\mathbf{q}|)$ and $\Phi^{(t)}(|\mathbf{q}|)$ depends on the relative momentum $q = (n - p')/2$ and contains full information about the structure of deuteron with one on-shell nucleon.

According to [2], the deuteron structure can be described by assuming the possible existence of non-nucleon or quark degrees of freedom. On the other hand, the shape of a high momentum tail of the nucleon distribution in the deuteron can be found from the true Regge asymptotic at $x \rightarrow 2$ [3], and the corresponding parameters can be fitted from

a good description of the inclusive proton spectrum in deuteron fragmentation $Dp \rightarrow pX$ [2, 3]. According to [2, 3], one can write the following form for $\Phi^{(u)}(|\mathbf{q}|)$:

$$\Phi^{(u)}(|\mathbf{q}|) = \frac{E_{\mathbf{k}}/E_{\mathbf{q}}}{2(1-x)} \tilde{\Phi}^{(u)}(|\mathbf{k}|) . \quad (4)$$

where

$$\tilde{\Phi}^{(u)}(|\mathbf{k}|) = (1 - \alpha_{2(3q)}) \left[U^2(|\mathbf{k}|) + W^2(|\mathbf{k}|) \right] + \alpha_{2(3q)} \frac{8\pi x(1-x)}{E_{\mathbf{k}}} G_{2(3q)}(x, \mathbf{k}_{\perp}) . \quad (5)$$

The parameter $\alpha_{2(3q)}$ is the probability for a non-nucleon component in the deuteron which is a state of two colorless ($3q$) systems.

$$G_{2(3q)}(x, \vec{k}_{\perp}) = \frac{b^2}{2\pi} \frac{\Gamma(A+B+2)}{\Gamma(A+1)\Gamma(B+1)} x^A (1-x)^B e^{-bk_{\perp}} . \quad (6)$$




Figure 1: The invariant pion spectrum [4] calculated within the relativistic impulse approximation with including of the non-nucleon component in the DWF [3, 2]; its probability $\alpha_{2(3q)}$ is $0.02 \div 0.04$ (dot-dashed and dashed curves, respectively).

including the non-nucleon degrees of freedom we have to require that the square of the new DWF has to be equal to the one determined by the Eq.(5). Introducing a mixing parameter $\alpha = (\pi/4)a$ one can find the forms of new $\mathcal{S}$ - and $\mathcal{D}$ -waves as the following:

$$\tilde{U}(|\mathbf{k}|) = \sqrt{1 - \alpha_{2(3q)}} U(|\mathbf{k}|) + \cos(\alpha) \Delta(|\mathbf{k}|) ; \quad (7)$$

$$\tilde{W}(|\mathbf{k}|) = \sqrt{1 - \alpha_{2(3q)}} W(|\mathbf{k}|) + \sin(\alpha) \Delta(|\mathbf{k}|) , \quad (8)$$

Fig. 1 presents the invariant pion spectrum calculated within the relativistic impulse approximation including the non-nucleon component in the DWF [2, 3]. One can see a good description of the experimental data [4] at all cumulative variable x_c .

However in some sense, the information contained in T_{20} from $Dp \rightarrow pX$ and $Dp \rightarrow \pi X$ processes is redundant, because the main ingredient by analyzing these reactions within the impulse approximation are the same deuteron properties. Therefore calculation of the tensor analyzing power including non-nucleon degrees of freedom in fragmentation of deuteron to pions can give us a new independent information concerning the deuteron structure at small $N - N$ distances. Actually, in [3] a form of $\tilde{\Phi}^{(u)}(|\mathbf{k}|)$ has been constructed only. However, to calculate T_{20} it is not enough, the corresponding orbital waves have to be known. Let us assume, the non-nucleon degrees of freedom result in a main contribution to the $\mathcal{S}$ - and $\mathcal{D}$ -waves of the DWF.

Constructing new forms of these waves by including the non-nucleon degrees of freedom we have to require that the square of the new DWF has to be equal to the one determined by the Eq.(5). Introducing a mixing parameter $\alpha = (\pi/4)a$ one can find the forms of new $\mathcal{S}$ - and $\mathcal{D}$ -waves as the following:

where the function Δ has been obtained from the equation:

$$\tilde{\Phi}^{(u)}(|\mathbf{k}|) = \tilde{U}^2(|\mathbf{k}|) + \tilde{W}^2(|\mathbf{k}|). \quad (9)$$




Figure 2: The tensor analyzing power T_{20} [5] calculated within the relativistic impulse approximation including of the non-nucleon component in the DWF [3, 2] with probability $\alpha_{2(3q)} = 0.03$. The solid and dashed lines represent the calculations using the mixing parameter $a = 0.0$ and 2.3 , respectively, Eqs. (7,8).

spectrum at all the values of x_c rather well, Fig. 1, and improve the description of data [5] concerning the analyzing power T_{20} in the fragmentation of deuteron to pions, Fig. 2.

Fig. 2 presents the analyzing power T_{20} calculated by using the functions $\tilde{U}, \tilde{W}$ including the non-nucleon components in the DWF, according to [2, 3]. It is shown from Fig. 2 the inclusion of non-nucleon components in the DWF improves the description of the experimental data about T_{20} at $x_c > 1$. The value of the parameter entering the Eqs.(7,8) $a = 2.3$ results in an optimal description of this observable.

Main results can be summarized as the following. Very interesting experimental data on T_{20} [5] showing approximately zero values at $x_c \geq 1$ are not reproduced by a theoretical calculus using even different kinds of the relativistic DWF [1]. This may indicate a possible existence of non-nucleon degrees of freedom or basically new mechanism of pion production in the kinematic region forbidden for free $N - N$ scattering.

The inclusion of the non-nucleon degrees of freedom within the approach suggested in [3, 2] allows us to describe experimental data about the inclusive pion

The support from RFFI grant N99-02-17727 is gratefully acknowledged by authors.

References

- [1] A.Yu. Illarionov, A.G. Litvinenko, G.I. Lykasov, Preprint JINR E1-2000-258, P2-2001-204; hep-ph/0012290, hep-ph/0007358.
- [2] G.I. Lykasov, EPAN **24**, 59 (1993).
- [3] A.V. Efremov *et al*, Sov. J. Nucl. Phys. **47**, 868 (1988).
- [4] A.M. Baldin, Nucl. Phys. **A434**, 695 (1985).
- [5] S.V. Afanasiev *et al*, Nucl. Phys. **A625**, 817 (1997); Phys. Lett. **B445**, 14 (1998); Preprint JINR E2-98-319.